

Table 1.11C Illicit Drug Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2009 and 2010

Age Category	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	971	1,026	600	646	457	488
12	31	27	26	23	20	16
13	32	37	24	31	16	22
14	42	38	34	32	24	24
15	57	51	47	44	28	31
16	55	54	48	47	34	36
17	60	62	53	57	38	40
18	88	87	76	77	54	58
19	92	86	77	73	57	57
20	89	91	72	76	56	60
21	97	91	76	72	59	54
22	96	83	77	66	55	52
23	86	84	67	61	46	49
24	91	83	65	60	49	47
25	78	87	52	62	40	45
26-29	334	345	220	221	152	169
30-34	355	352	193	209	146	172
35-39	352	348	169	169	128	121
40-44	396	375	182	175	129	129
45-49	390	419	164	182	117	136
50-54	563	567	226	240	180	196
55-59	443	538	184	218	147	132
60-64	381	365	135	134	108	97
65 or Older	355	369	119	135	103	97

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.11D Illicit Drug Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2009 and 2010

Age Category	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.39	0.40	0.24	0.25	0.18	0.19
12	0.75	0.69	0.66	0.58	0.52	0.41
13	0.76	0.81	0.60	0.69	0.40	0.51
14	0.86	0.87	0.72	0.76	0.55	0.60
15	1.00	0.98	0.88	0.91	0.61	0.70
16	1.01	1.07	0.94	0.95	0.74	0.79
17	1.03	1.07	0.98	1.05	0.78	0.84
18	1.16	1.16	1.10	1.15	0.91	1.02
19	1.21	1.25	1.21	1.23	1.03	1.08
20	1.23	1.31	1.29	1.32	1.14	1.14
21	1.25	1.34	1.23	1.25	1.18	1.08
22	1.24	1.19	1.30	1.19	1.09	1.05
23	1.19	1.30	1.29	1.20	1.00	1.07
24	1.22	1.31	1.27	1.23	1.09	1.04
25	1.37	1.32	1.15	1.28	0.98	1.05
26-29	1.21	1.22	1.09	1.10	0.82	0.92
30-34	1.09	1.12	0.90	0.93	0.71	0.80
35-39	1.21	1.18	0.80	0.80	0.62	0.60
40-44	1.16	1.15	0.78	0.80	0.58	0.60
45-49	1.04	1.13	0.67	0.75	0.51	0.59
50-54	1.58	1.63	0.96	1.06	0.78	0.87
55-59	1.65	1.80	0.95	1.06	0.77	0.66
60-64	1.81	1.87	0.85	0.80	0.69	0.58
65 or Older	0.93	0.95	0.31	0.35	0.27	0.25

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.11P Illicit Drug Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2010 Versus 2009

Age Category	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.9724		0.5062		0.3272	
12	0.6392		0.9367		0.4926	
13	0.8028		0.1951		0.0503	
14	0.3652		0.9449		0.8093	
15	0.8926		1.0000		0.6256	
16	0.0538		0.2110		0.8744	
17	0.7278		0.3768		0.9782	
18	0.5120		0.5987		0.5751	
19	0.6882		0.9716		0.0373	
20	0.1134		0.2386		0.3863	
21	0.1821		0.1647		0.8172	
22	0.0334		0.1298		0.8024	
23	0.5920		0.7187		0.4635	
24	0.4219		0.3439		0.5935	
25	0.6495		0.6166		0.9109	
26-29	0.3388		0.9261		0.7788	
30-34	0.6697		0.1755		0.0203	
35-39	0.7161		0.6230		0.8823	
40-44	0.9747		0.8815		0.6847	
45-49	0.5545		0.4068		0.3812	
50-54	0.8477		0.4776		0.7957	
55-59	0.1808		0.5446		0.1830	
60-64	0.4698		0.7042		0.6104	
65 or Older	0.3762		0.4423		0.5668	

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.12C Marijuana Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2009 and 2010

Age Category	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	953	1,021	541	557	395	406
12	10	9	9	9	7	4
13	18	21	15	19	9	11
14	30	28	27	26	20	18
15	42	42	38	39	25	28
16	47	47	43	41	31	33
17	55	56	46	53	36	38
18	83	81	71	71	51	54
19	86	81	72	67	55	53
20	82	87	67	73	51	58
21	93	86	72	66	53	49
22	91	80	70	61	50	48
23	82	82	56	57	42	44
24	88	77	62	52	44	42
25	72	82	48	55	36	40
26-29	314	328	198	199	131	151
30-34	336	336	172	186	128	146
35-39	331	322	140	140	101	102
40-44	377	366	138	143	97	110
45-49	370	404	133	151	101	116
50-54	543	545	191	187	146	159
55-59	418	517	146	182	122	119
60-64	356	352	113	106	92	87
65 or Older	313	347	96	113	92	67

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.12D Marijuana Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2009 and 2010

Age Category	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.38	0.40	0.21	0.22	0.16	0.16
12	0.28	0.23	0.25	0.22	0.20	0.11
13	0.46	0.48	0.38	0.44	0.24	0.27
14	0.65	0.68	0.60	0.63	0.46	0.45
15	0.81	0.86	0.77	0.82	0.53	0.62
16	0.94	0.96	0.88	0.86	0.69	0.73
17	1.00	1.02	0.88	1.01	0.73	0.80
18	1.17	1.17	1.08	1.14	0.89	0.97
19	1.20	1.26	1.17	1.20	1.01	1.03
20	1.26	1.30	1.29	1.28	1.08	1.12
21	1.31	1.32	1.23	1.23	1.10	1.03
22	1.29	1.23	1.24	1.15	1.01	1.02
23	1.21	1.34	1.15	1.17	0.93	0.99
24	1.28	1.30	1.26	1.11	0.99	0.95
25	1.45	1.36	1.14	1.21	0.91	0.94
26-29	1.24	1.24	1.03	1.02	0.72	0.84
30-34	1.10	1.14	0.82	0.85	0.63	0.69
35-39	1.22	1.15	0.68	0.67	0.50	0.51
40-44	1.16	1.18	0.65	0.67	0.46	0.52
45-49	1.06	1.15	0.55	0.64	0.44	0.51
50-54	1.60	1.66	0.83	0.84	0.64	0.71
55-59	1.62	1.86	0.76	0.90	0.64	0.60
60-64	1.79	1.83	0.72	0.64	0.59	0.52
65 or Older	0.82	0.89	0.25	0.29	0.24	0.17

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.12P Marijuana Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2010 Versus 2009

Age Category	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.4465		0.5246		0.3338	
12	0.3454		0.9670		0.3631	
13	0.3746		0.1274		0.4125	
14	0.8769		0.7307		0.7688	
15	0.7481		0.5811		0.5114	
16	0.8991		0.8240		0.7273	
17	0.4788		0.1063		0.7709	
18	0.7128		0.5396		0.3595	
19	0.5871		0.7152		0.1768	
20	0.1017		0.3520		0.5998	
21	0.3827		0.2591		0.6620	
22	0.0686		0.3016		0.9190	
23	0.6232		0.7784		0.3557	
24	0.1540		0.3961		0.9588	
25	0.3921		0.8397		0.7040	
26-29	0.2012		0.7529		1.0000	
30-34	0.8864		0.0717		0.0517	
35-39	0.9388		0.6961		0.4992	
40-44	0.9953		0.9520		0.2993	
45-49	0.7118		0.5920		0.7013	
50-54	0.9915		0.7803		0.9699	
55-59	0.1116		0.4647		0.3725	
60-64	0.8437		0.9953		0.8433	
65 or Older	0.3992		0.4352		0.8112	

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.13C Cocaine Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2009 and 2010

Age Category	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	655	682	211	203	136	108
12	4	0	4	0	4	*
13	2	5	2	1	0	*
14	5	6	4	6	2	1
15	10	11	8	11	4	6
16	14	15	11	13	6	7
17	20	17	16	13	9	6
18	28	29	21	21	10	14
19	37	31	26	23	13	12
20	37	40	22	23	11	13
21	48	42	31	28	15	15
22	50	42	30	29	16	12
23	41	45	20	23	12	16
24	48	45	23	25	12	13
25	42	47	19	24	8	14
26-29	168	184	80	91	38	38
30-34	189	202	80	85	49	55
35-39	196	192	54	68	29	34
40-44	208	227	60	62	42	34
45-49	240	274	65	51	47	37
50-54	338	372	94	62	65	41
55-59	258	295	41	45	31	4
60-64	182	180	40	29	37	17
65 or Older	163	164	12	*	*	*

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.13D Cocaine Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2009 and 2010

Age Category	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.26	0.27	0.08	0.08	0.05	0.04
12	0.12	0.01	0.10	0.01	0.10	*
13	0.06	0.13	0.06	0.03	0.01	*
14	0.13	0.15	0.10	0.14	0.05	0.02
15	0.22	0.27	0.18	0.26	0.09	0.14
16	0.33	0.36	0.25	0.32	0.15	0.18
17	0.46	0.40	0.37	0.30	0.21	0.14
18	0.54	0.58	0.42	0.42	0.19	0.29
19	0.76	0.67	0.55	0.51	0.29	0.28
20	0.84	0.81	0.51	0.50	0.26	0.27
21	1.05	0.94	0.72	0.65	0.35	0.36
22	1.06	0.93	0.70	0.68	0.40	0.29
23	0.94	1.02	0.49	0.56	0.29	0.39
24	1.04	1.02	0.53	0.61	0.29	0.31
25	1.07	1.10	0.52	0.60	0.23	0.35
26-29	0.92	0.95	0.46	0.53	0.22	0.23
30-34	0.86	0.89	0.40	0.42	0.25	0.27
35-39	0.91	0.89	0.27	0.35	0.15	0.18
40-44	0.87	0.97	0.28	0.30	0.20	0.17
45-49	0.89	1.04	0.28	0.23	0.21	0.17
50-54	1.32	1.46	0.42	0.28	0.29	0.19
55-59	1.26	1.34	0.22	0.23	0.17	0.02
60-64	1.13	1.05	0.26	0.18	0.24	0.10
65 or Older	0.43	0.42	0.03	*	*	*

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.13P Cocaine Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2010 Versus 2009

Age Category	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.7060		0.2565		0.3024	
12	0.0326		0.0795		*	
13	0.3513		0.6169		*	
14	0.6632		0.7041		0.3090	
15	0.3133		0.1896		0.1793	
16	0.7984		0.5272		0.6874	
17	0.1197		0.3965		0.2524	
18	0.3785		0.6232		0.5381	
19	0.0357		0.4430		0.9965	
20	0.0633		0.2097		0.9621	
21	0.0505		0.0660		0.7886	
22	0.1963		0.2795		0.3134	
23	0.9122		0.6976		0.8095	
24	0.1322		0.8424		0.8418	
25	0.4979		0.7339		0.1599	
26-29	0.6937		0.9810		0.8974	
30-34	0.2826		0.5105		0.5995	
35-39	0.7539		0.6767		0.8710	
40-44	0.9031		0.7417		0.4413	
45-49	0.2124		0.6677		0.4824	
50-54	0.8506		0.1818		0.5578	
55-59	0.4484		0.9162		0.0570	
60-64	0.9133		0.5789		0.3270	
65 or Older	0.8968		*		*	

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.14C Crack Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2009 and 2010

Age Category	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	316	378	106	91	77	60
12	2	0	2	0	1	*
13	1	5	1	0	0	*
14	2	3	2	1	2	*
15	4	4	4	3	3	2
16	6	2	3	2	0	0
17	5	8	3	3	1	1
18	10	13	7	11	2	9
19	12	12	6	10	4	7
20	14	16	7	4	3	*
21	20	17	7	5	2	3
22	16	18	6	7	0	6
23	19	18	5	9	3	8
24	21	16	8	5	8	2
25	18	19	9	8	3	3
26-29	80	84	33	31	24	8
30-34	100	105	29	15	25	4
35-39	106	99	29	33	17	18
40-44	111	127	31	41	26	31
45-49	121	149	40	33	33	26
50-54	164	197	65	38	42	35
55-59	106	127	18	4	18	4
60-64	61	79	24	26	24	13
65 or Older	24	34	*	*	*	*

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.14D Crack Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2009 and 2010

Age Category	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.13	0.15	0.04	0.04	0.03	0.02
12	0.04	0.01	0.04	0.01	0.04	*
13	0.03	0.12	0.03	0.01	0.01	*
14	0.06	0.07	0.06	0.02	0.04	*
15	0.10	0.11	0.09	0.06	0.07	0.04
16	0.15	0.06	0.06	0.04	0.01	0.01
17	0.11	0.19	0.07	0.08	0.02	0.02
18	0.20	0.27	0.14	0.22	0.03	0.18
19	0.25	0.26	0.13	0.22	0.10	0.17
20	0.33	0.34	0.17	0.08	0.08	*
21	0.48	0.40	0.18	0.11	0.04	0.07
22	0.40	0.42	0.15	0.18	0.01	0.14
23	0.46	0.45	0.13	0.21	0.06	0.20
24	0.49	0.39	0.19	0.13	0.19	0.06
25	0.50	0.48	0.24	0.20	0.08	0.07
26-29	0.47	0.49	0.20	0.19	0.14	0.05
30-34	0.50	0.51	0.15	0.07	0.13	0.02
35-39	0.51	0.50	0.14	0.17	0.09	0.09
40-44	0.52	0.59	0.15	0.20	0.12	0.15
45-49	0.52	0.65	0.18	0.15	0.15	0.12
50-54	0.72	0.86	0.29	0.17	0.19	0.16
55-59	0.56	0.65	0.10	0.02	0.10	0.02
60-64	0.39	0.48	0.16	0.16	0.16	0.08
65 or Older	0.06	0.09	*	*	*	*

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.14P Crack Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: P Values from Tests of Differences of Percentages, 2010 Versus 2009

Age Category	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.1444		0.2790		0.2397	
12	0.3277		0.2908		*	
13	0.5410		0.2765		*	
14	0.5349		0.2344		*	
15	0.9467		0.5659		0.6444	
16	0.0216		0.4464		0.9326	
17	0.0915		0.8128		0.7602	
18	0.5582		0.5278		0.3670	
19	0.8397		0.1235		0.4271	
20	0.4466		0.0882		*	
21	0.2313		0.0696		0.3053	
22	0.5167		0.2266		0.0077	
23	0.5488		0.6706		0.2003	
24	0.0574		0.6381		0.4024	
25	0.3860		0.3500		0.8902	
26-29	0.8215		0.8580		0.2342	
30-34	0.2498		0.1331		0.0808	
35-39	0.6366		0.9088		0.8074	
40-44	0.2339		0.6847		0.8858	
45-49	0.0112		0.6610		0.9908	
50-54	0.9595		0.2409		0.9521	
55-59	0.7229		0.0911		0.1749	
60-64	0.3931		0.8337		0.4201	
65 or Older	0.6213		*		*	

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.15C Hallucinogen Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2009 and 2010

Age Category	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	665	689	158	177	91	84
12	8	4	2	3	2	0
13	11	10	9	9	3	6
14	16	14	15	13	8	8
15	19	18	16	17	9	10
16	20	23	18	20	10	9
17	28	28	24	25	12	12
18	38	38	32	32	18	19
19	45	42	35	31	19	16
20	46	49	31	33	15	16
21	51	51	36	33	14	19
22	61	49	38	33	22	13
23	49	52	20	28	7	13
24	51	50	21	28	9	17
25	47	52	20	21	9	10
26-29	206	202	69	73	46	39
30-34	234	221	75	73	54	34
35-39	186	203	45	30	22	6
40-44	182	196	36	28	*	18
45-49	211	240	20	27	3	20
50-54	325	331	21	56	21	17
55-59	252	298	13	*	*	*
60-64	174	187	*	10	*	*
65 or Older	110	124	25	10	*	*

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.15D Hallucinogen Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2009 and 2010

Age Category	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.26	0.27	0.06	0.07	0.04	0.03
12	0.23	0.11	0.07	0.08	0.04	0.01
13	0.28	0.25	0.22	0.22	0.09	0.16
14	0.38	0.35	0.35	0.32	0.20	0.22
15	0.43	0.41	0.36	0.39	0.20	0.25
16	0.46	0.54	0.42	0.48	0.23	0.22
17	0.62	0.64	0.52	0.58	0.28	0.27
18	0.69	0.72	0.61	0.63	0.36	0.39
19	0.90	0.87	0.72	0.66	0.40	0.36
20	0.99	0.96	0.70	0.68	0.37	0.33
21	1.11	1.09	0.83	0.74	0.34	0.44
22	1.19	1.01	0.85	0.74	0.52	0.31
23	1.02	1.13	0.47	0.68	0.16	0.32
24	1.05	1.09	0.49	0.66	0.23	0.41
25	1.12	1.17	0.54	0.52	0.26	0.26
26-29	1.06	1.03	0.40	0.43	0.27	0.24
30-34	1.02	0.95	0.38	0.36	0.28	0.17
35-39	0.85	0.89	0.23	0.16	0.11	0.03
40-44	0.77	0.86	0.17	0.14	*	0.09
45-49	0.84	0.96	0.09	0.12	0.01	0.09
50-54	1.29	1.33	0.09	0.26	0.09	0.08
55-59	1.22	1.40	0.07	*	*	*
60-64	1.08	1.09	*	0.06	*	*
65 or Older	0.29	0.32	0.07	0.03	*	*

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.15P Hallucinogen Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: P Values from Tests of Differences of Percentages, 2010 Versus 2009

Age Category	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.9606		0.9194		0.6274	
12	0.2365		0.6712		0.1268	
13	0.4355		0.7680		0.4482	
14	0.8392		0.9949		0.1739	
15	0.6654		0.4465		0.3722	
16	0.3661		0.7364		0.1631	
17	0.5559		0.7865		0.6553	
18	0.5163		0.5328		0.9357	
19	0.9014		0.9188		0.6901	
20	0.1896		0.2198		0.6553	
21	1.0000		0.6147		0.2852	
22	0.1878		0.8461		0.1363	
23	0.9748		0.1181		0.0315	
24	0.8034		0.0802		0.0594	
25	0.7947		0.4022		0.7301	
26-29	0.5687		0.9279		0.9349	
30-34	0.8380		0.7329		0.1057	
35-39	0.2485		0.1004		0.0472	
40-44	0.2818		0.9745		*	
45-49	0.9237		0.3718		0.3890	
50-54	0.2501		0.1780		0.7508	
55-59	0.3741		*		*	
60-64	0.8107		*		*	
65 or Older	0.9054		0.5774		*	

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.16C Inhalant Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2009 and 2010

Age Category	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	473	473	109	114	57	82
12	22	23	17	16	12	8
13	25	27	17	21	9	14
14	29	25	18	18	7	10
15	31	31	17	18	7	9
16	28	26	19	16	9	7
17	28	24	14	14	8	8
18	32	30	17	19	6	7
19	35	32	20	16	11	6
20	31	35	12	18	7	5
21	41	36	20	14	8	7
22	39	35	16	13	9	8
23	35	37	17	18	4	12
24	36	32	10	13	4	9
25	34	36	8	6	6	3
26-29	142	140	43	33	19	15
30-34	183	173	21	32	9	20
35-39	156	159	27	24	26	14
40-44	153	157	19	24	8	14
45-49	164	176	30	20	19	19
50-54	206	214	54	74	28	65
55-59	154	188	24	3	*	3
60-64	128	106	*	18	*	18
65 or Older	98	73	*	19	*	*

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.16D Inhalant Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2009 and 2010

Age Category	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.19	0.19	0.04	0.04	0.02	0.03
12	0.57	0.58	0.44	0.42	0.34	0.21
13	0.62	0.64	0.44	0.50	0.24	0.34
14	0.67	0.62	0.43	0.47	0.17	0.26
15	0.66	0.71	0.38	0.42	0.17	0.22
16	0.63	0.61	0.43	0.38	0.21	0.17
17	0.61	0.54	0.31	0.32	0.18	0.19
18	0.61	0.59	0.34	0.38	0.12	0.15
19	0.71	0.70	0.44	0.36	0.24	0.13
20	0.69	0.72	0.28	0.38	0.17	0.12
21	0.94	0.77	0.49	0.31	0.19	0.16
22	0.87	0.79	0.39	0.31	0.23	0.18
23	0.82	0.86	0.42	0.44	0.10	0.29
24	0.81	0.76	0.24	0.31	0.11	0.22
25	0.86	0.86	0.21	0.14	0.17	0.09
26-29	0.79	0.77	0.26	0.20	0.11	0.09
30-34	0.85	0.79	0.11	0.16	0.04	0.10
35-39	0.74	0.76	0.14	0.12	0.13	0.07
40-44	0.70	0.72	0.09	0.12	0.04	0.07
45-49	0.68	0.75	0.13	0.09	0.08	0.09
50-54	0.89	0.93	0.24	0.34	0.13	0.29
55-59	0.80	0.93	0.13	0.02	*	0.02
60-64	0.81	0.63	*	0.11	*	0.11
65 or Older	0.26	0.19	*	0.05	*	*

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.16P Inhalant Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2010 Versus 2009

Age Category	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.1639		0.6370		0.2311	
12	0.6246		0.3424		0.1502	
13	0.9322		0.5535		0.1356	
14	0.0450		0.8912		0.0806	
15	0.5641		0.5876		0.1491	
16	0.0696		0.0288		0.1703	
17	0.1362		0.4161		0.6087	
18	0.8646		0.9805		0.8638	
19	0.9661		0.8648		0.3557	
20	0.9808		0.7619		0.1124	
21	0.0993		0.1095		0.9034	
22	0.4843		0.8697		0.5741	
23	0.6275		0.8616		0.3842	
24	0.0232		0.7453		0.6943	
25	0.5381		0.4617		0.1368	
26-29	0.7527		0.3704		0.3713	
30-34	0.5933		0.1104		0.0714	
35-39	0.9863		0.9636		0.7828	
40-44	0.9051		0.1954		0.1792	
45-49	0.8055		0.2395		0.8388	
50-54	0.7790		0.7257		0.3639	
55-59	0.4445		0.3900		*	
60-64	0.2173		*		*	
65 or Older	0.4804		*		*	

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.17C Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2009 and 2010

Age Category	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	738	785	367	398	250	268
12	19	19	17	16	12	14
13	19	22	14	18	10	13
14	25	23	21	19	15	14
15	35	30	29	25	18	16
16	36	35	29	31	17	17
17	40	39	32	33	17	17
18	54	53	41	44	27	28
19	60	55	46	41	26	26
20	57	61	42	47	27	29
21	63	62	48	45	30	28
22	72	61	49	43	32	27
23	62	57	40	37	22	24
24	58	61	38	41	25	24
25	54	66	34	38	22	23
26-29	234	228	144	130	90	79
30-34	234	234	115	126	73	93
35-39	220	216	114	104	85	65
40-44	234	233	110	116	84	75
45-49	256	261	101	108	59	65
50-54	333	385	137	178	97	127
55-59	267	317	124	130	91	57
60-64	226	207	80	83	56	43
65 or Older	223	226	66	89	48	71

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.17D Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2009 and 2010

Age Category	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.29	0.31	0.15	0.16	0.10	0.11
12	0.49	0.49	0.44	0.41	0.31	0.36
13	0.47	0.52	0.37	0.44	0.25	0.31
14	0.57	0.56	0.49	0.46	0.35	0.36
15	0.70	0.66	0.60	0.57	0.40	0.39
16	0.78	0.77	0.64	0.68	0.40	0.41
17	0.83	0.84	0.68	0.72	0.39	0.39
18	0.92	0.92	0.75	0.81	0.51	0.55
19	1.10	1.07	0.89	0.86	0.55	0.57
20	1.10	1.16	0.88	0.95	0.62	0.62
21	1.24	1.17	1.03	0.90	0.69	0.61
22	1.31	1.15	1.02	0.93	0.73	0.62
23	1.21	1.18	0.89	0.85	0.52	0.58
24	1.15	1.24	0.85	0.95	0.58	0.56
25	1.22	1.27	0.86	0.88	0.60	0.58
26-29	1.12	1.12	0.79	0.73	0.52	0.47
30-34	1.01	0.99	0.58	0.61	0.38	0.46
35-39	0.97	0.96	0.56	0.52	0.42	0.34
40-44	0.94	1.01	0.50	0.56	0.38	0.36
45-49	0.96	0.99	0.43	0.47	0.26	0.29
50-54	1.33	1.50	0.60	0.79	0.43	0.57
55-59	1.31	1.42	0.66	0.66	0.49	0.30
60-64	1.34	1.18	0.51	0.50	0.36	0.26
65 or Older	0.59	0.58	0.17	0.23	0.13	0.18

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.17P Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2010 Versus 2009

Age Category	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.6509		0.8698		0.9220	
12	0.4856		0.6752		0.4935	
13	0.7739		0.3521		0.2660	
14	0.2223		0.4190		0.8056	
15	0.7041		0.4136		0.3215	
16	0.0518		0.6226		0.7229	
17	0.4476		0.9114		0.2875	
18	0.3831		0.5508		0.5020	
19	0.7960		0.9324		0.5055	
20	0.2424		0.1321		0.3218	
21	0.5471		0.3621		0.9308	
22	0.1220		0.7148		0.1852	
23	0.1302		0.1742		0.5203	
24	0.9901		0.8649		0.3990	
25	0.0086		0.6486		0.8096	
26-29	0.9887		0.9425		0.7701	
30-34	0.6923		0.7774		0.0469	
35-39	0.4345		0.7288		0.2704	
40-44	0.6864		0.8938		0.7105	
45-49	0.2571		0.9779		0.4123	
50-54	0.4223		0.1448		0.2698	
55-59	0.2986		0.8029		0.1477	
60-64	0.1448		0.3052		0.4623	
65 or Older	0.8780		0.3048		0.1851	

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.18C Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2009 and 2010

Age Category	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	583	608	322	336	214	214
12	18	18	16	16	11	13
13	18	21	14	17	9	12
14	22	22	20	17	13	13
15	31	29	27	24	17	15
16	33	33	26	28	16	15
17	38	37	30	31	16	16
18	50	47	37	37	25	25
19	54	50	41	38	23	22
20	51	55	36	41	23	26
21	56	52	41	38	26	23
22	58	54	40	39	26	24
23	56	51	35	31	17	20
24	53	56	33	35	21	19
25	49	59	29	33	20	22
26-29	206	200	126	121	69	68
30-34	205	205	100	123	65	88
35-39	179	190	101	88	78	58
40-44	187	189	95	94	77	64
45-49	192	185	89	95	54	60
50-54	261	261	114	142	76	102
55-59	203	211	111	116	78	39
60-64	141	161	58	76	44	29
65 or Older	113	130	52	56	43	47

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.18D Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2009 and 2010

Age Category	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.23	0.24	0.13	0.13	0.09	0.08
12	0.47	0.47	0.42	0.40	0.30	0.35
13	0.46	0.49	0.35	0.41	0.24	0.29
14	0.50	0.54	0.46	0.43	0.30	0.34
15	0.64	0.64	0.57	0.55	0.38	0.36
16	0.73	0.74	0.59	0.64	0.38	0.35
17	0.79	0.78	0.64	0.67	0.36	0.38
18	0.87	0.86	0.68	0.72	0.48	0.50
19	1.03	1.01	0.82	0.81	0.48	0.49
20	1.02	1.06	0.78	0.84	0.53	0.56
21	1.19	1.05	0.90	0.79	0.60	0.52
22	1.12	1.09	0.85	0.86	0.59	0.56
23	1.15	1.11	0.80	0.74	0.42	0.50
24	1.08	1.19	0.77	0.82	0.50	0.46
25	1.15	1.19	0.76	0.78	0.53	0.54
26-29	1.03	1.04	0.71	0.69	0.41	0.40
30-34	0.92	0.90	0.51	0.59	0.33	0.44
35-39	0.83	0.87	0.50	0.45	0.39	0.30
40-44	0.79	0.85	0.43	0.45	0.35	0.31
45-49	0.79	0.76	0.39	0.42	0.24	0.27
50-54	1.10	1.10	0.50	0.64	0.34	0.46
55-59	1.04	1.03	0.59	0.59	0.41	0.20
60-64	0.88	0.94	0.37	0.46	0.29	0.18
65 or Older	0.30	0.33	0.14	0.14	0.11	0.12

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.18P Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2010 Versus 2009

Age Category	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.5360		0.5388		0.5266	
12	0.3825		0.4139		0.2875	
13	0.7824		0.2713		0.5551	
14	0.4170		0.1465		0.7557	
15	0.8099		0.3679		0.3926	
16	0.2905		0.9661		0.2696	
17	0.2086		0.3720		0.3896	
18	0.1889		0.3335		0.4495	
19	0.6645		0.8317		0.7293	
20	0.3655		0.2181		0.2933	
21	0.3852		0.3006		0.8144	
22	0.1724		0.9454		0.8196	
23	0.2094		0.0426		0.7233	
24	0.8068		0.6645		0.5715	
25	0.0077		0.9344		0.8337	
26-29	0.8974		0.9326		0.9293	
30-34	0.9167		0.4163		0.1702	
35-39	0.0670		0.8663		0.4147	
40-44	0.6669		0.9602		0.5980	
45-49	0.3765		0.9175		0.2160	
50-54	0.1707		0.6674		0.6510	
55-59	0.4877		0.6468		0.1050	
60-64	0.7894		0.8066		0.2297	
65 or Older	0.2699		0.9160		0.7757	

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.